

Third order integrability conditions for homogeneous potentials of degree -1

Thierry Combet¹ and Christoph Koutschan²

¹*IMCCE (France)*^{a)}

²*RISC, Johannes Kepler University Linz (Austria)*^{b)}

(Dated: 30 October 2012)

We prove an integrability criterion of order 3 for a homogeneous potential of degree -1 in the plane. Still, this criterion depends on some integer and it is impossible to apply it directly except for families of potentials whose eigenvalues are bounded. To address this issue, we use holonomic and asymptotic computations with error control of this criterion and apply it to the potential of the form $V(r, \theta) = r^{-1}h(\exp(i\theta))$ with $h \in \mathbb{C}[z]$, $\deg h \leq 3$. We then find all meromorphically integrable potentials of this form.

PACS numbers: 02.30.Hq 02.70.Wz

Keywords: Non-integrability, Homogeneous potentials, Differential Galois theory, Holonomicity

^{a)}Electronic mail: combot@imcce.fr

^{b)}Electronic mail: ckoutsch@risc.jku.at; ¹ Supported by University Paris 7 Diderot.; ² Partially supported by the DDMF project of the MSR-INRIA Joint Centre, and by the Austrian Science Fund (FWF): P20162-N18.

I. INTRODUCTION

In this article, we will be interested in non-integrability proofs of meromorphic homogeneous potentials of degree -1 in the plane, and in particular in nongeneric cases. Writing our potential V in polar coordinates, and making the Fourier expansion in the angle gives us

$$V(r, \theta) = r^{-1} \sum_{k=-\infty}^{\infty} a_k e^{ik\theta}. \quad (1)$$

This type of potential covers many physical problems in celestial mechanics and n -body problems, in particular the anisotropic Kepler problem, the isosceles 3-body problem, the colinear 3-body problem, the symmetric 4-body problem and so on. Moreover, for such a potential there are strong integrability conditions, thanks to the Morales-Ramis theory¹ and to a very effective criterion of Yoshida². Still, for such a general potential, this criterion will not be sufficient. This is not particularly because this class of potentials is large, but because there are nongeneric, very resistant cases inside. For example, if we want to study the integrability of $V(r, \theta) = r^{-1}h(\exp(i\theta))$ with a polynomial h , we have a priori a potential with $\deg h + 1$ complex parameters, and Yoshida's integrability criterion will restrict this family to a family with $\deg h - 1$ integer parameters. Still one would like to have a finite list of possible integrable potentials, so as to be able to check the existence of first integrals one by one. Here we will present a stronger criterion in Theorems 2 and 3 which is able to deal with such families, and which therefore is capable to settle any integrability question on finite dimensional families of type (1). As an application of our method, we will apply this criterion in the case $V(r, \theta) = r^{-1}h(\exp(i\theta))$ with $h \in \mathbb{C}[z]$, $\deg h \leq 3$. To do precise statements, let us now begin with some definitions concerning homogeneous potentials and integrability.

Definition 1. *We consider the algebraic variety $\mathcal{S} = \{(q_1, q_2, r) \in \mathbb{C}^3, r^2 = q_1^2 + q_2^2\}$ and the derivations for a function f on \mathcal{S}*

$$\frac{\partial f}{\partial q_1} = \partial_1 f + r^{-1} q_1 \partial_3 f, \quad \frac{\partial f}{\partial q_2} = \partial_2 f + r^{-1} q_2 \partial_3 f$$

where ∂_i is the derivative according to the i -th variable (the variables of f are q_1, q_2, r in this order). This defines a symplectic form on $\mathbb{C}^2 \times \mathcal{S}$ on which we consider a Hamiltonian H of the form

$$H(p_1, p_2, q_1, q_2, r) = \frac{1}{2}(p_1^2 + p_2^2) + V(q_1, q_2, r)$$

Integrability conditions for homogeneous potentials

with the associated system of differential equations

$$\dot{r} = r^{-1}(q_1\dot{q}_1 + q_2\dot{q}_2), \quad \dot{q}_i = \frac{\partial}{\partial p_i}H, \quad \dot{p}_i = \frac{\partial}{\partial q_i}H, \quad i = 1, 2. \quad (2)$$

The potential V is assumed to be meromorphic on \mathcal{S} and to have the following form in polar coordinates:

$$V(r, \theta) = \frac{1}{r}U(\theta), \quad r \cos \theta = q_1, \quad r \sin \theta = q_2$$

This implies that V is homogeneous of degree -1 . We say that I is a meromorphic first integral of H , if I is a meromorphic function on $\mathbb{C}^2 \times \mathcal{S}$ such that

$$\dot{I} = \{H, I\} = \sum_{i=1}^2 \left(\frac{\partial}{\partial p_i}H \frac{\partial}{\partial q_i}I - \frac{\partial}{\partial q_i}H \frac{\partial}{\partial p_i}I \right) = 0.$$

Obviously, the Hamiltonian H itself is a first integral. We will say that V is meromorphically integrable if it possesses an additional meromorphic first integral which is independent almost everywhere from H .

Definition 2. We call $c = (c_1, c_2, c_3) \in \mathcal{S}$ a Darboux point of V if

$$\frac{\partial}{\partial q_1}V(c) = \alpha c_1 \quad \text{and} \quad \frac{\partial}{\partial q_2}V(c) = \alpha c_2 \quad (3)$$

where $\alpha \in \mathbb{C}$ is called the multiplier. Because V has singularities, we will **always** assume that $c_3 \neq 0$. Because of homogeneity, we can always choose $\alpha = 0$ or $\alpha = -1$. We say that c is non-degenerate if $\alpha \neq 0$. To the Darboux point c we associate a homothetic orbit given by

$$r(t) = c_3\phi(t), \quad q_i(t) = c_i\phi(t), \quad p_i(t) = c_i\dot{\phi}(t) \quad (i = 1, 2), \quad (4)$$

with ϕ satisfying the following differential equation

$$\frac{1}{2}\dot{\phi}(t)^2 = -\frac{\alpha}{\phi(t)} + E, \quad E \in \mathbb{C}.$$

In the following, we will often omit the last component of a Darboux point $c \in \mathcal{S}$ as it is defined up to a sign (and the choice of sign does not matter) by the two first components.

Definition 3. The first order variational equation of H near a homothetic orbit is given by

$$\ddot{X}(t) = \frac{1}{\phi(t)^3} \nabla^2 V(c) X(t)$$

Integrability conditions for homogeneous potentials

where $\nabla^2 V(c)$ is the Hessian of V (according to derivations in q). After diagonalization (if possible) and the change of variable $\phi(t) \rightarrow t$, the equation simplifies to

$$2t^2(Et + 1)\ddot{X}_i - t\dot{X}_i = \lambda_i X_i,$$

where the λ_i are the eigenvalues of the Hessian of V evaluated at the Darboux point c , i.e., $\lambda_i \in \text{Sp}(\nabla^2 V(c))$.

Theorem 1. (Morales, Ramis, Yoshida²³¹⁴) *If V is meromorphically integrable, then the neutral component of the Galois group of the variational equation near a homothetic orbit with $E \neq 0$ is abelian at all orders. If we fix the multiplier of the associated Darboux point to -1 , the Galois group of the first order variational equation has an abelian neutral component if and only if*

$$\text{Sp}(\nabla^2 V(c)) \subset \left\{ \frac{1}{2}(k-1)(k+2) : k \in \mathbb{N} \right\}.$$

If the multiplier of the Darboux point is 0, the Galois group of the first order variational equation has an abelian neutral component if and only if

$$\text{Sp}(\nabla^2 V(c)) \subset \{0\}.$$

In fact, this is not exactly the same statement as the original theorem because we allow r to appear in the potential and in the first integrals.

Proof. Let $\Gamma \subset \mathbb{C}^2 \times \mathcal{S}$ denote the curve defined by equation (4) without the singular point $(q_1, q_2, r) = (0, 0, 0)$, and M an open neighbourhood of Γ in $\mathbb{C}^2 \times \mathcal{S}$ such that H is holomorphic on M . The Hamiltonian H is then well defined and holomorphic on a symplectic manifold M and the additional first integral is meromorphic on M . Hence, using the main theorem of³, the neutral component of the Galois group of the variational equation near Γ is abelian at all orders over the base field of meromorphic functions on Γ . The variational equation is a hypergeometric equation. In⁶, Kimura classifies Galois groups of hypergeometric equations over the base field $\mathbb{C}(t)$. We can use this classification as the Galois group over the base field $\mathbb{C}(t)$ is the same as over the base field of meromorphic functions because the hypergeometric equation is a Fuchsian equation (see page 73 of¹). This produces the condition on the spectrum of $\nabla^2 V(c)$. The case of a degenerate Darboux point leads to the variational equation

$$\ddot{X} = \lambda t^{-3} X$$

which is a Bessel equation (after a change of variables). Its Galois group over the field of meromorphic functions in t has not an abelian identity component except if $\lambda = 0$. \square

Note that in the case of a degenerate Darboux point, we explicitly need that the first integral is meromorphic including $r = 0$, as the variational equation is not regular singular at this point.

II. MAIN RESULTS

In this section, we are going to state the main theorems of this article. The remaining parts of this paper are dedicated to their proofs.

Theorem 2. *Let V be a homogeneous potential of degree -1 in the plane. We suppose that $c = (1, 0)$ is a Darboux point of V with multiplier -1 . If the variational equation is integrable at order 3, then the following conditions are fulfilled*

$$\mathrm{Sp}(\nabla^2 V(c)) = \left\{ 2, \frac{1}{2}(p-1)(p+2) \right\} \text{ for some } p \in \mathbb{N}.$$

If p is even then

$$\left(\frac{\partial^3 V}{\partial q_1 \partial q_2^2} \right)^2 f_1(p) + \left(\frac{\partial^3 V}{\partial q_2^3} \right)^2 f_2(p) + \left(\frac{\partial^4 V}{\partial q_2^4} \right) f_3(p) = 0,$$

and if p is odd then

$$\frac{\partial^3 V}{\partial q_2^3} = 0 \quad \text{and} \quad \left(\frac{\partial^3 V}{\partial q_1 \partial q_2^2} \right)^2 f_1(p) + \left(\frac{\partial^4 V}{\partial q_2^4} \right) f_3(p) = 0,$$

where the functions f_1, f_2, f_3 satisfy explicit P -finite recurrences, i.e., linear recurrences with polynomial coefficients.

This theorem is a generalization of the criterion given by Yoshida for homogeneous potentials in the case of degree -1 and dimension 2. A similar theorem could be proven in higher dimensions, but the main problem is that Theorem 2 is almost inapplicable in this form. In most cases, it is necessary to study more closely the expression of the functions $f_1(p), f_2(p), f_3(p)$ to apply it, and for the moment, because of limitations of computing power, it seems only possible to do in dimension 2 (for which the computations are already tedious).

Theorem 3. *The functions $f_1(2n), f_2(2n), f_3(2n)$ can be written as*

$$\begin{aligned} f_1(2n) &= \epsilon_1(n) \left(\frac{1511011}{67108864n^2} - \frac{1511011}{134217728n^3} + \frac{31731231}{4294967296n^4} \right) \\ f_2(2n) &= \epsilon_2(n) \left(\frac{22665165}{1073741824n^4} - \frac{22665165}{1073741824n^5} + \frac{298125}{4194304n^6} \right) \\ f_3(2n) &= \epsilon_3(n) \left(-\frac{1740684681}{68719476736n^2} + \frac{1740684681}{137438953472n^3} - \frac{2400813907}{68719476736n^4} \right) \end{aligned} \quad (5)$$

with

$$|\epsilon_i(n) - 1| \leq 10^{-5} \quad \forall n \geq 100.$$

With this, we can apply Theorem 2 to some concrete examples:

Theorem 4. *Let V be a potential in the plane expressed in polar coordinates by*

$$V(r, \theta) = r^{-1} \left(a + be^{i\theta} + ce^{2i\theta} + de^{3i\theta} \right). \quad (6)$$

If V is meromorphically integrable, then V belongs to one of the following families

$$\begin{aligned} V = r^{-1}a, \quad V = r^{-1}(a + be^{i\theta}), \quad V = r^{-1}(ae^{i\theta} + be^{3i\theta}), \\ V = r^{-1}(a + be^{2i\theta}), \quad V = r^{-1}(a + be^{3i\theta}), \quad V = r^{-1}(a + be^{i\theta})^3, \end{aligned} \quad (7)$$

with $a, b \in \mathbb{C}$.

The first three families are integrable, with a polynomial first integral of degree 1 or 2 in p . The status of the last three families is unknown. This is not due to an incomplete application of the Morales-Ramis Theorem, but linked to the fact that either they do not possess any Darboux points, or in the last case the only Darboux point is very degenerate and therefore the Morales-Ramis Theorem gives no integrability constraints at any order, as proven in⁵.

In practical problems like Theorem 4, studying integrability only using the Morales-Ramis criterion is impossible because of two facts. First we need a Darboux point of our problem; if we do not have any, the only thing we can do is to try to find an additional first integral using the direct method of Hietarinta⁷.

The second problem is the following scenario: inside the family of potentials given by Theorem 4, there exist submanifolds in the space of parameters for which the potential possesses only one Darboux point and the eigenvalue at this Darboux point can be arbitrarily high. In this case, the higher variational method is required. But the constraint at order 2 does not give sufficient conditions to conclude, and it is necessary to go to order 3.

But the expression of this constraint cannot be written explicitly for all possible eigenvalues, only for a finite number of them. To apply this third-order criterion, we derive P-finite recurrences and asymptotic expansions with error control in Theorem 3. This allows us to prove that the integrability condition is not fulfilled. The proof of Theorem 4 therefore will be split into two parts:

1. The first part consists in constructing a manifold M in the space of the parameters a, b, c, d such that if the eigenvalues for all Darboux points are real, then the parameters belong to M . Then we produce a decomposition $M = M_1 \cup \dots \cup M_k$ and study each manifold separately. For some of them, the corresponding potentials possess sufficiently many Darboux points to give a strong enough condition for integrability only using the Morales-Ramis criterion at order 1 (there could exist some resistant cases for which a higher variational equation is needed but without the phenomenon of arbitrary high eigenvalues like in⁸). But for specific cases, this phenomenon occurs. It has already been noticed by Maciejewski in⁹ who lets this specific case open.
2. The second part will be devoted to these specific manifolds M_i where the Morales-Ramis criterion at order 1 is almost powerless. We use Theorems 2 and 3 to solve these hard cases.

In¹⁹, the authors deal with a similar difficulty with the spring pendulum for which there is a discrete infinite set of parameters for which there are no obstructions to integrability at order 2. They also study third order variational equations, but then use analytic tools to study a sequence of monodromy elements, and finally prove that this sequence never vanishes. Thanks to our explicit expression via P -finite recurrences, such a problem can be analysed more systematically here.

III. EIGENVALUE BOUNDING

Definition 4. *We will denote*

$$\mathcal{M} = \left\{ V(r, \theta) = r^{-1}U(\theta) \text{ with } U \text{ meromorphic and } 2\pi\text{-periodic} \right\}.$$

Integrability conditions for homogeneous potentials

Let $V \in \mathcal{M}$. We denote by $d(V)$ the set of Darboux points c of V with multiplier -1 and $c_3 \neq 0$. For $c \in d(V)$ we have $\text{Sp}(\nabla^2 V(c)) = \{2, \lambda\}$ and we denote

$$\Lambda(c) = \begin{cases} \lambda & \text{if } \lambda \in \mathbb{R} \\ -\infty & \text{otherwise} \end{cases}.$$

Definition 5. We consider a subset $E \subset \mathcal{M}$ and define

$$\Lambda(E) = \sup_{V \in E, d(V) \neq \emptyset} \inf_{c \in d(V)} \Lambda(c).$$

We say that the problem of finding all meromorphically integrable potentials in E is a bounded eigenvalue problem if $\Lambda(E) < \infty$.

Remark 1. We have $\Lambda(\mathcal{M}) = \infty$ because of the following family

$$V(r, \theta) = r^{-1} \left((1 + a) - 2ae^{i\theta} + ae^{2i\theta} \right), \quad a \in \mathbb{R},$$

for which only one Darboux point $c = (1, 0)$ exists; the corresponding eigenvalue is $\lambda = 2a - 1$. This proves that the family of potentials considered in Theorem 4 is an unbounded eigenvalue problem.

Lemma 1. For a potential $V \in \mathcal{M}$ the Darboux points c such that $c_3 \neq 0$ can be written as $c = (c_1, c_2) = (r_0 \cos(\theta_0), r_0 \sin(\theta_0))$ with θ_0 being a critical point of U . The Darboux point c is non-degenerate if and only if $U(\theta_0) \neq 0$ and in this case, the eigenvalues of the Hessian of V , evaluated at c , are

$$\text{Sp}(\nabla^2 V(c)) = \left\{ 2, \frac{U''(\theta_0)}{U(\theta_0)} - 1 \right\}.$$

if we choose the multiplier of c to be -1 .

Proof. For $V = r^{-1}U(\theta)$ the conditions (3) that c is a Darboux point are:

$$\begin{aligned} r_0^{-3} (-c_1 U(\theta_0) - c_2 U'(\theta_0)) &= \alpha c_1, \\ r_0^{-3} (-c_2 U(\theta_0) + c_1 U'(\theta_0)) &= \alpha c_2. \end{aligned}$$

Assuming $c_3 \neq 0$, it follows that $U(\theta_0) = -\alpha r_0^3$ and $U'(\theta_0) = 0$, which means that θ_0 is a critical point of U . Since $c_3 \neq 0$ implies that $r_0 \neq 0$, we see that the case $\alpha = 0$ (degenerate

Darboux point) is equivalent to $U(\theta_0) = 0$. Setting $\alpha = -1$ and $U'(\theta_0) = 0$ we get the Hessian matrix

$$\nabla^2 V(c) = \frac{1}{r_0^3} \begin{pmatrix} (2c_1^2 - c_2^2)U(\theta_0) + c_2^2 U''(\theta_0) & c_1 c_2 (3U(\theta_0) - U''(\theta_0)) \\ c_1 c_2 (3U(\theta_0) - U''(\theta_0)) & (2c_2^2 - c_1^2)U(\theta_0) + c_1^2 U''(\theta_0) \end{pmatrix}$$

whose eigenvalues are exactly those claimed above (using $U(\theta_0) = r_0^3$). \square

Recall that the potentials given by (6) are $V(r, \theta) = r^{-1}U(\theta)$ with $U(\theta) = a + be^{i\theta} + ce^{2i\theta} + de^{3i\theta}$. We now assume that V possesses at least one non-degenerate Darboux point c with $c_3 \neq 0$. After rotation, we can always assume that $c = (1, 0)$ is a Darboux point. As shown in Lemma 1, it corresponds to a critical point for $\theta = 0$. Moreover, because this Darboux point is non-degenerate, we know that $U(0) \neq 0$. Then by dilatation, we can also suppose that $U(0) = 1$ and get the following equations

$$U(0) = a + b + c + d = 1,$$

$$U'(0) = i(b + 2c + 3d) = 0.$$

Solving these equations for c and d , yields the expression

$$V_{a,b} = r^{-1} \left(a + be^{i\theta} + (3 - 3a - 2b)e^{2i\theta} + (2a + b - 2)e^{3i\theta} \right)$$

for the potentials where $a, b \in \mathbb{C}$.

Theorem 5. *If $V_{a,b}$ is meromorphically integrable, then it belongs to one of the following families*

$$\begin{aligned} E_1 &= r^{-1} \left(-\frac{1}{3}b + 1 + be^{i\theta} - be^{2i\theta} + \frac{1}{3}be^{3i\theta} \right), \\ E_2 &= r^{-1} \left(-\frac{1}{6}k(k+1)e^{3i\theta} + \frac{1}{4}k(k+1)e^{2i\theta} - \frac{1}{12}k^2 - \frac{1}{12}k + 1 \right), \\ E_3 &= r^{-1} \left(-\frac{1}{4}k(k+1)e^{2i\theta} + \frac{1}{2}k(k+1)e^{i\theta} - \frac{1}{4}k^2 - \frac{1}{4}k + 1 \right), \\ E_4 &= r^{-1} \left(\frac{(s - 6\lambda_2)\lambda_2}{18(\lambda_1 + \lambda_2)} e^{3i\theta} - \frac{(3\lambda_1 + s - 3\lambda_2)\lambda_2}{6(\lambda_1 + \lambda_2)} e^{2i\theta} + \right. \\ &\quad \left. \frac{(6\lambda_1 + s)\lambda_2}{6(\lambda_1 + \lambda_2)} e^{i\theta} + \frac{-9\lambda_1\lambda_2 - \lambda_2s + 18\lambda_1 + 18\lambda_2 - 3\lambda_2^2}{18(\lambda_1 + \lambda_2)} \right) \end{aligned}$$

where $b \in \mathbb{C}$ and $k \in \mathbb{N}$. The quantities arising in E_4 are

$$s^2 = 6\lambda_1^2\lambda_2 + 6\lambda_1\lambda_2^2 - 36\lambda_1\lambda_2,$$

$$\lambda_i = \frac{1}{2}(k_i - 1)(k_i + 2) + 1 \quad (i = 1, 2),$$

with $k_1 \in \mathbb{N} \setminus \{0, 3\}$ and $k_2 \in \mathbb{N}^*$.

Proof. For all non-degenerate Darboux points $c = (\gamma \cos(\theta_0), \gamma \sin(\theta_0))$ the corresponding eigenvalue λ satisfies

$$U''(\theta_0) - (\lambda + 1)U(\theta_0) = 0 \quad \text{and} \quad U'(\theta_0) = 0 \quad (8)$$

(note that this condition is also satisfied if c is degenerate). We write $U(\theta) = h_{a,b}(\exp(i\theta))$, $U'(\theta) = izh'_{a,b}(\exp(i\theta))$, and $U''(\theta) = \tilde{h}_{a,b}(\exp(i\theta))$ with

$$\begin{aligned} h_{a,b}(z) &= a + bz + (3 - 3a - 2b)z^2 + (2a + b - 2)z^3, \\ \tilde{h}_{a,b}(z) &= -bz - 4(3 - 3a - 2b)z^2 - 9(2a + b - 2)z^3. \end{aligned}$$

So to find the eigenvalues of all Darboux points, one just needs to compute the following resultant which corresponds to the conditions (8):

$$\begin{aligned} P_{a,b}(\lambda) &= \text{res}_z(\tilde{h}_{a,b}(z) - (\lambda + 1)h_{a,b}(z), h'_{a,b}(z)) \\ &= (2a + b - 2)(6a + 2b - 6 + (\lambda + 1))(-18ab^2 - 6b^3 + 18b^2 + (\lambda + 1) \\ &\quad \times (108a^3 + 108a^2b - 216a^2 + 36ab^2 + 108a - 108ab - 9b^2 + 4b^3)) \end{aligned}$$

All the roots of $P_{a,b}(\lambda)$ correspond to an eigenvalue of some Darboux point, except possibly in those cases (a, b) where $P_{a,b}$ vanishes as a polynomial in λ or in the case where $h'_{a,b}(z)$ has the root 0.

Let us begin with the special cases. We compute the points $(a, b) \in \mathbb{C}^2$ for which $P_{a,b} = 0$ in $\mathbb{C}[\lambda]$. We find that it is the zero set of the ideal $\langle 2a + b - 2 \rangle \cap \langle a, b \rangle$. Moreover, the polynomial $h'_{a,b}(z)$ has a zero root if and only if $b = 0$. So, all the specific cases belong to the zero set of $\langle 2a + b - 2 \rangle \cap \langle b \rangle$. First, for $b = 0$ we find

$$\begin{aligned} Q_1 &= \text{res}_z(\tilde{h}_{a,0}(z) - (\lambda + 1)h_{a,0}(z), h'_{a,0}(z)/z, z) \\ &= 216(a - 1)^3(6a - 6 + (\lambda + 1)), \end{aligned}$$

and second, for $b = 2 - 2a$ we get

$$\begin{aligned} Q_2 &= \text{res}_z(\tilde{h}_{a,2-2a}(z) - (\lambda + 1)h_{a,2-2a}(z), h'_{a,2-2a}(z), z) \\ &= -4(a - 1)^2(2a - 2 + (\lambda + 1)). \end{aligned}$$

As we know that the eigenvalues should be of the form $\frac{1}{2}(k - 1)(k + 2)$, $k \in \mathbb{N}$, we obtain the potentials E_2 and E_3 from these two cases.

Now for the generic case, we express a and b depending on the roots of $P_{a,b}(\lambda)$ and obtain the expression E_4 . Since it is not valid for $k_1 = k_2 = 0$, we study this case separately and find the condition $a = -\frac{1}{3}b + 1$, which gives E_1 . Note that fixing $\lambda_1 = 0$ in E_4 yields the potential E_2 , whereas $\lambda_1 = 6$ results in E_3 . The case $k_2 = 0$ produces $V = r^{-1}$ which already belongs to E_1 . \square

Corollary 1. *With the same notation as in Theorem 5, we have $\Lambda(E_1) = -1$ and $\Lambda(E_2) = \Lambda(E_3) = \Lambda(E_4) = \infty$.*

Remark 2. *The types of E_2 , E_3 and E_4 differ fundamentally although they are all unbounded eigenvalue problems. This is because the dimension of E_4 is 2 and the dimension of E_2 and E_3 is only 1. Because of that, we could call E_4 a doubly unbounded eigenvalue problem because it possesses two Darboux points whose eigenvalues can be independently arbitrarily high. Because of that, we will need to apply a third order integrability criterion simultaneously at the two Darboux points. The potential E_1 has only one Darboux point with eigenvalue -1 . This eigenvalue belongs to the Morales-Ramis table and so higher variational methods will be required, but only for this fixed eigenvalue (which is much easier).*

In the parameter space, we get 4 algebraic manifolds. For E_2 , E_3 , and E_4 , a tedious treatment with higher variational equations is required. For E_1 we will be able to check integrability easily with Theorem 2. A similar procedure could be applied to any set of homogeneous potentials depending rationally on some parameters. Here computing power is the main limitation; in particular, because for typical problems, the number of parameters is much smaller than the number of roots which requires resultant computations and prime ideal decompositions. One should note that we have deliberately chosen a set of potentials (6) which is particularly difficult to treat. For most common problems (outside the general complete classification), these unbounded eigenvalue manifolds have small dimension (1 in the case found by⁹) or even inexistent like in⁸ or¹⁰.

IV. HIGHER ORDER VARIATIONAL METHODS

We will first recall some properties of the solutions of the first order variational equations. After diagonalisation and in the integrable case, the equation is the following (after fixing

Integrability conditions for homogeneous potentials

the energy $E = 1$)

$$2t^2(1+t)y''(t) - ty'(t) - \frac{1}{2}(n-1)(n+2)y(t) = 0 \quad (n \in \mathbb{N}). \quad (9)$$

After the change of variables $t \rightarrow (t^2 - 1)^{-1}$, this equation becomes

$$(t^2 - 1)y''(t) + 4ty'(t) - (n-1)(n+2)y(t) = 0 \quad (n \in \mathbb{N}). \quad (10)$$

A basis of solutions is given by (P_n, Q_n) where P_n are polynomials in t (for $n \geq 1$) and the functions Q_n are

$$Q_n(t) = P_n(t) \int \frac{1}{(t^2 - 1)^2 P_n(t)^2} dt.$$

The functions Q_n are multivalued except for $n = 0$ which will be a special case. Indeed, the Galois group of (9) in this case is Id instead of \mathbb{C} .

The polynomials P_n can be computed using the Rodrigues type formula

$$P_n(t) = \frac{1}{t^2 - 1} \frac{\partial^{n-1}}{\partial t^{n-1}} (t^2 - 1)^n \quad (n \geq 1) \quad (11)$$

which also gives a normalisation for their leading coefficient. The functions Q_n can be written as

$$Q_n(t) = \epsilon_n P_n(t) \operatorname{arctanh}\left(\frac{1}{t}\right) + \frac{W_n(t)}{t^2 - 1} \quad (n \geq 1) \quad (12)$$

where W_n are polynomials given by

$$W_{2k}(t) = \frac{(-1)^k (t^2 - 1)}{2^{4k}} \left(\frac{\pi {}_2F_1\left(\frac{1}{2} - k, k + 1, \frac{1}{2}, t^2\right)}{\Gamma\left(k + \frac{1}{2}\right)^2} + \frac{2kt(2k + 1) \operatorname{arctanh}(t) {}_2F_1\left(1 - k, k + \frac{3}{2}, \frac{3}{2}, t^2\right)}{(k!)^2} \right),$$

$$W_{2k+1}(t) = \frac{(-1)^k (t^2 - 1)}{2^{4k+2}} \left(\frac{\pi t(k + 1)(2k + 1) {}_2F_1\left(\frac{1}{2} - k, k + 2, \frac{3}{2}, t^2\right)}{\Gamma\left(k + \frac{3}{2}\right)^2} - \frac{2 \operatorname{arctanh}(t) {}_2F_1\left(-k, k + \frac{3}{2}, \frac{1}{2}, t^2\right)}{(k!)^2} \right)$$

and ϵ_n is a real sequence given by

$$\epsilon_n = \frac{n(n+1)}{4^n (n!)^2}.$$

Conventionally, we will take for $n = 0$:

$$P_0(t) = \frac{t}{t^2 - 1}, \quad Q_0(t) = \frac{1}{t^2 - 1}.$$

Lemma 2. *The functions $P_n(t)$ and $\frac{1}{e_n}Q_n(t)$ satisfy the differential equation (10) and the three-term recurrence*

$$4n(n+1)(n+2)y_n(t) - 2t(n+2)(2n+3)y_{n+1}(t) + (n+3)y_{n+2}(t) = 0.$$

Proof. Given the explicit expressions (11) and (12) we can use holonomic closure properties to derive the differential equation resp. recurrence they satisfy. We first express (11) as

$$P_n(t) = \frac{(n-1)!}{2\pi i(t^2-1)} \oint \frac{(u^2-1)^n}{(u-t)^n} du$$

by Cauchy's differentiation formula. By the method of creative telescoping we obtain the differential equation and the recurrence (this calculation was carried out by the software package `HolonomicFunctions`^{11,12}). Similarly we can apply holonomic closure properties to the closed form expression (12). \square

Lemma 3. *(proved in¹³) Let $F \in \mathbb{C}(z_1)[z_2]$ and $f(t) = F\left(t, \operatorname{arctanh}\left(\frac{1}{t}\right)\right)$. We consider the field extension*

$$K = \mathbb{C}\left(t, \operatorname{arctanh}\left(\frac{1}{t}\right), \int f dt\right)$$

and the monodromy group $G = \sigma(K, \mathbb{C}(t))$. If G is abelian, then

$$\frac{\partial}{\partial \alpha} \operatorname{Res}_{t=\infty} F\left(t, \operatorname{arctanh}\left(\frac{1}{t}\right) + \alpha\right) = 0.$$

Proof. We will consider two paths, the ‘‘eight’’ path σ_1 around the singularities -1 and 1 , and the path σ_2 around infinity. At infinity, the function $F\left(t, \operatorname{arctanh}\left(\frac{1}{t}\right) + \alpha\right)$ will have a series expansion of the kind

$$\int F\left(t, \operatorname{arctanh}\left(\frac{1}{t}\right) + \alpha\right) dt = \sum_{n=n_0}^{\infty} a_n(\alpha)t^n + r(\alpha) \ln t$$

because the function $\operatorname{arctanh}\left(\frac{1}{t}\right)$ has a regular point at infinity. Let us now consider the monodromy commutator

$$\sigma = \sigma_2^{-1} \sigma_1^{-\frac{\beta}{2i\pi}} \sigma_2 \sigma_1^{\frac{\beta}{2i\pi}}.$$

We have that $\sigma_1^{\frac{\beta}{2i\pi}}(f) = F\left(t, \operatorname{arctanh}\left(\frac{1}{t}\right) + \beta\right)$ and $\sigma_2(\ln t) = \ln t + 2i\pi$. We deduce that

$$\sigma(f) = f + r(\beta) - r(0).$$

This $r(\alpha)$ corresponds to the residue of $F\left(t, \operatorname{arctanh}\left(\frac{1}{t}\right) + \alpha\right)$ at infinity. If the monodromy is commutative, then the commutator σ should act trivially on f . This is the case only if $r(\beta) - r(0) = 0$ for all $\beta \in \mathbb{Z}$. The function r is a polynomial in β , so $r(\beta) - r(0) = 0$ for all $\beta \in \mathbb{C}$. From this the claim follows. \square

Integrability conditions for homogeneous potentials

In the following, we will also need to use the next lemma which is a kind of reciprocal version of Lemma 3.

Lemma 4. (proved in¹³) We consider

$$F(t) = \sum_{i=0}^3 H_i(t) \operatorname{arctanh} \left(\frac{1}{t} \right)^i$$

with $H_0, \dots, H_3 \in \mathbb{C}[t]$. If the conditions of Lemma 3 are satisfied, then

- If $\operatorname{Res}_{t=\infty} F(t) = 0$, then $\int F dt \in \mathbb{C} \left[t, \operatorname{arctanh} \left(\frac{1}{t} \right) \right]$
- If $\operatorname{Res}_{t=\infty} F(t) \neq 0$, then $\int F dt \in \mathbb{C} \left[t, \operatorname{arctanh} \left(\frac{1}{t} \right), \ln(t^2 - 1) \right]$

Theorem 6. Let V be a homogeneous potential of degree -1 in the plane. We suppose that $c = (1, 0)$ is a Darboux point of V with multiplier -1 . If the variational equation is integrable at order 2 then

$$\operatorname{Sp} \left(\nabla^2 V(c) \right) = \left\{ 2, \frac{1}{2}(p-1)(p+2) \right\}, \quad p \in \mathbb{N},$$

and for odd p we have $\frac{\partial^3 V}{\partial q_2^3} = 0$.

This theorem is in fact a particular case of Theorem 2 in¹³ for which the three indices i, j, k are equal.

Remark 3. Because the constraint appears only for odd p , the variational equations of order 2 give no constraint for even p . Hence this is not sufficient for proving non-integrability for an unbounded manifold.

V. PROOF OF THEOREM 2

Proof. The variational equation at order 3 is given by

$$\begin{aligned}
\ddot{X}_1 &= \frac{2}{\phi^3}X_1 + \frac{1}{2}\frac{a}{\phi^4}Y_{1,1} - \frac{4b}{3\phi^5}Z^3 \\
\ddot{X}_2 &= \frac{\lambda}{\phi^3}X_2 + \frac{a}{\phi^4}Y_{2,1} + \frac{b}{\phi^4}Y_{1,1} + \frac{c}{\phi^5}Z^3 \\
\dot{Y}_{1,1} &= 2Y_{1,2} \\
\dot{Y}_{1,2} &= \frac{\lambda}{\phi^3}Y_{1,1} + \frac{b}{\phi^4}Z^3 + Y_{1,3} \\
\dot{Y}_{1,3} &= \frac{\lambda}{\phi^3}Y_{1,2} + \frac{b}{\phi^4}Z^2\dot{Z} \\
\dot{Y}_{2,1} &= Y_{2,2} + Y_{2,3} \\
\dot{Y}_{2,2} &= \frac{2}{\phi^3}Y_{2,1} - \frac{4b}{3\phi^5}Z^3 + Y_{2,4} \\
\dot{Y}_{2,3} &= \frac{\lambda}{\phi^3}Y_{2,1} + Y_{2,4} \\
\dot{Y}_{2,4} &= \frac{2}{\phi^3}Y_{2,3} - \frac{4b}{3\phi^5}Z^2\dot{Z} + \frac{\lambda}{\phi^3}Y_{2,2} \\
\ddot{Z} &= \frac{2}{\phi^3}Z
\end{aligned}$$

where $\lambda = \frac{1}{2}(n-1)(n+2)$. The coefficients a, b, c correspond to the following derivatives

$$a = \frac{\partial^3}{\partial q_1 \partial q_2^2} V(c), \quad b = \frac{1}{2} \frac{\partial^3}{\partial q_3^2} V(c), \quad c = \frac{1}{6} \frac{\partial^4}{\partial q_2^4} V(c),$$

and the others are given using the Euler relation for homogeneous functions. A complete procedure to build these equations is given by¹⁴. The functions $Y_{1,1}$ and $Y_{2,1}$ are solutions of a system of linear differential equations with an inhomogeneous term, and the homogeneous part is in fact a symmetric product of the first order variational equation. Here, we already put to zero terms that we think in advance they will not produce integrability constraints. As before, we use the change of variables $\phi(t) \longrightarrow (t^2 - 1)^{-1}$.

We choose $Z(t) = Q_n$ and compute the solution for X_2 of the above system. We first remark that X_2 is in the Picard-Vessiot field, so it is also the case for its derivative. We now perform integration by parts and see that one term is already in the Picard-Vessiot field, and the other is

$$\int 2(t^2 - 1)^2 \left(a^2 t P_n Q_n I_1 + 4b^2 P_n^2 Q_n I_2 + c(t^2 - 1) Q_n^4 \right) dt \quad (13)$$

where

$$\begin{aligned}
 I_1 &= \int \left(\frac{\int \left(\frac{t(t^2-1)^2 Q_n^3}{P_n} + \frac{I_3}{(t^2-1)^2 P_n^2} \right) dt}{t^2(t^2-1)^2} + \frac{\int \frac{I_3}{t^2(t^2-1)^2} dt}{(t^2-1)^2 P_n^2} \right) dt \\
 I_2 &= \int \frac{\int \left((t^2-1)^2 Q_n^3 + \frac{2}{(t^2-1)^2 P_n^2} \int (t^2-1)^4 P_n Q_n^2 (P_n \dot{Q}_n - Q_n \dot{P}_n) dt \right) dt}{(t^2-1)^2 P_n^2} dt \\
 I_3 &= \int t(t^2-1)^4 Q_n^2 (P_n \dot{Q}_n - Q_n \dot{P}_n) dt
 \end{aligned}$$

Let us now study this expression term by term. We begin with the third summand of (13) which is

$$2c \int (t^2 - 1)^3 Q_n^4 dt.$$

It has already the form of Lemma 3. So as in the proof of Lemma 3, the monodromy commutator will be computed using

$$\text{Res}_{t=\infty} (t^2 - 1)^3 (Q_n + \epsilon_n \alpha P_n)^4 dt.$$

Now look at the term in b^2 . It is not as complicated as we could think because of the following relation

$$P_n \dot{Q}_n - \dot{P}_n Q_n = (t^2 - 1)^{-2} \quad \forall n \in \mathbb{N}$$

which is linked to the Wronskian of Equation (10). Thanks to that, the term in b^2 can be written as

$$8b^2 \int P_n^2 Q_n (t^2 - 1)^2 \int \frac{\int (t^2 - 1)^2 Q_n^3 + 2 \frac{\int P_n Q_n^2 (t^2 - 1)^2 dt}{(t^2 - 1)^2 P_n^2} dt}{(t^2 - 1)^2 P_n^2} dt dt$$

and then using integration by parts, this gives

$$16b^2 \int Q_n^3 (t^2 - 1)^2 \int P_n Q_n^2 (t^2 - 1)^2 dt dt - 8b^2 \int P_n Q_n^2 (t^2 - 1)^2 dt \int Q_n^3 (t^2 - 1)^2 dt$$

Now by Lemma 4 we have for all even integers $n > 1$:

$$\int P_n Q_n^2 (t^2 - 1)^2 dt, \int Q_n^3 (t^2 - 1)^2 dt \in \mathbb{C}(t) \left[\text{arctanh} \left(\frac{1}{t} \right) \right].$$

So we are integrating a polynomial in arctanh with rational coefficients, and this corresponds to the hypotheses of Lemma 3. The second term does not provide any monodromy, so we only have to study the first term and thus the sequence

$$\text{Res}_{t=\infty} (Q_n + \epsilon_n \alpha P_n)^3 (t^2 - 1)^2 \int P_n (Q_n + \epsilon_n \alpha P_n)^2 (t^2 - 1)^2 dt.$$

Integrability conditions for homogeneous potentials

Now we look at the term in a^2 . It can be simplified to

$$\int 2a^2(t^2 - 1)^2 P_n Q_n t \int \frac{\int \frac{(t^2-1)^2 Q_n^3 t}{P_n} + \frac{\int (t^2-1)^2 Q_n^2 t dt}{(t^2-1)^2 P_n^2} dt}{t^2(t^2 - 1)^2} + \frac{\int \frac{(t^2-1)^2 Q_n^2 t dt}{t^2(t^2-1)^2} dt}{(t^2 - 1)^2 P_n^2} dt$$

We now use again integrations by parts (recall that $P_2 = 4t$):

$$8a^2 \int (t^2 - 1)^2 Q_n^2 Q_2 \int (t^2 - 1)^2 Q_n^2 t dt - 8a^2 \int (t^2 - 1)^2 Q_n^2 t \int (t^2 - 1)^2 Q_n^2 Q_2 dt.$$

To conclude we can again use Lemmas 3 and 4. We first prove that

$$\forall n \neq 1 \quad \int P_2 Q_n^2 (t^2 - 1)^2 dt, \int Q_n^2 Q_2 (t^2 - 1)^2 dt \in \mathbb{C}(t) \left[\operatorname{arctanh} \left(\frac{1}{t} \right) \right].$$

The case $n = 1$ corresponds to $\lambda = 0$, for which we have always the coefficient $a = 0$. Now we make a final integration by parts which gives

$$16a^2 \int (t^2 - 1)^2 Q_n^2 Q_2 \int (t^2 - 1)^2 Q_n^2 t dt - 8a^2 \int (t^2 - 1)^2 Q_n^2 t \int (t^2 - 1)^2 Q_n^2 Q_2 dt.$$

Thanks to that, we get a constraint of the form given by Theorem 2 and the coefficients are given by (multiplying them by ϵ_n^{-2} for further simplifications)

$$f_1(n) = \langle \alpha^3 \rangle 2\epsilon_n^{-2} \operatorname{Res}_{t=\infty} \left((t^2 - 1)^2 (Q_n + \epsilon_n \alpha P_n)^2 (Q_2 + \epsilon_2 \alpha P_2) \right. \\ \left. \times \int (t^2 - 1)^2 (Q_n + \epsilon_n \alpha P_n)^2 P_2 dt \right), \quad (14)$$

$$f_2(n) = \langle \alpha^3 \rangle 2\epsilon_n^{-2} \operatorname{Res}_{t=\infty} \left((t^2 - 1)^2 (Q_n + \epsilon_n \alpha P_n)^3 \right. \\ \left. \times \int (t^2 - 1)^2 (Q_n + \epsilon_n \alpha P_n)^2 P_n dt \right), \quad (15)$$

$$f_3(n) = \langle \alpha^3 \rangle \frac{1}{6} \epsilon_n^{-2} \operatorname{Res}_{t=\infty} \left((t^2 - 1)^3 (Q_n + \epsilon_n \alpha P_n)^4 \right), \quad (16)$$

where $\langle \cdot \rangle$ denotes coefficient extraction. In fact, only the coefficient of α^3 appears in these residues. We need not to prove this fact, because we simply select the coefficient of α^3 , ignoring the question whether the other coefficients are zero or not.

We now look at the case $n = 0$. All our previous calculations are also valid in this case except those involving Lemma 4 because we only have

$$\int P_0 Q_0^2 (t^2 - 1)^2 dt, \int Q_0^3 (t^2 - 1)^2 dt \in \mathbb{C}(t) \left[\operatorname{arctanh} \left(\frac{1}{t} \right), \ln(t^2 - 1) \right], \\ \int P_2 Q_0^2 (t^2 - 1)^2 dt, \int Q_0^2 Q_2 (t^2 - 1)^2 dt \in \mathbb{C}(t) \left[\operatorname{arctanh} \left(\frac{1}{t} \right) \right].$$

Integrability conditions for homogeneous potentials

So, the coefficients in a^2, c are also

$$2 \operatorname{Res}_{t=\infty} \left((t^2 - 1)^2 Q_0^2 Q_2 \int (t^2 - 1)^2 P_2 Q_0^2 dt \right),$$

$$\frac{1}{6} \operatorname{Res}_{t=\infty} \left((t^2 - 1)^3 Q_0^4 \right).$$

We find that these residues are both 0, and so the corresponding integral does not provide any additional monodromy. The case of the coefficient in b^2 is a little more difficult because the integral does not satisfy the conditions of Lemma 3. After an explicit computation, we arrive at the following integral

$$\int \frac{1}{t^2 - 1} \left(-t \operatorname{arctanh} \left(\frac{1}{t} \right) - \frac{1}{2} \ln(t^2 - 1) \right) dt =$$

$$\frac{1}{2} \ln(2) \ln(t - 1) + \frac{1}{2} \operatorname{dilog}(t + 1) +$$

$$\frac{1}{8} \ln(t + 1)^2 + \frac{1}{4} \ln(t + 1) \ln(t - 1) - \frac{1}{8} \ln(t - 1)^2.$$

All the terms are in $\mathbb{C}[t, \operatorname{arctanh}(\frac{1}{t}), \ln(t^2 - 1)]$ except one, namely the dilogarithmic term

$$\operatorname{dilog}(t + 1) = \int \frac{\ln(t + 1)}{t} dt.$$

With the same idea as in Lemma 3, we see that this term has a noncommutative monodromy because of the following residue in 0

$$\operatorname{Res}_{t=0} \frac{\ln(t + 1) + \alpha}{t} = \alpha$$

which depends explicitly on α . So, for $n = 0$, the integrability condition at order 3 is in fact just $b^2 = 0$.

□

VI. HOLONOMICITY AND ASYMPTOTICS

In this section we are going to derive P-finite recurrences (i.e., linear recurrences with polynomial coefficients) for the sequences $f_1(n)$, $f_2(n)$, and $f_3(n)$ that appeared in the previous section. The methods that we employ are based on Zeilberger's holonomic systems approach¹⁵. The recurrences presented below were computed with the method of creative telescoping, to which a brief introduction is given below (see¹¹ for more details).

Integrability conditions for homogeneous potentials

Let S_n denote the forward shift operator in n , i.e., $S_n f(n) = f(n+1)$, and D_x the derivative w.r.t. x , i.e., $D_x f(x) = f'(x)$. The method works for the class of holonomic functions, which in short are (multivariate) functions that are solutions of maximally overdetermined systems of linear difference and differential equations with polynomial coefficients. The set of all equations which a given holonomic function satisfies forms a left ideal (we call it *annihilating ideal*) in some Ore algebra of the form

$$\mathbb{C}(m, n, \dots, x, y, \dots) \langle S_m, S_n, \dots, D_x, D_y \dots \rangle.$$

The nice fact about holonomic functions is that this class is closed under certain operations (addition, multiplication, certain substitutions, definite summation and integration) which can be executed algorithmically: given the defining systems of equations for two holonomic functions f and g , there are algorithms to compute a holonomic system for $f + g$, $f \cdot g$, etc.

For computing integrals (or residues), the method of creative telescoping makes use of the fundamental theorem of calculus. Consider a definite integral of the form $\int_a^b f dx$ where the integrand f depends also on some other (discrete and/or continuous) parameters. We need f to be holonomic, i.e., there is some left ideal I of annihilating operators in the corresponding Ore algebra \mathbb{O} . The idea is now to come up with an operator $A + D_x B \in I$ where $A, B \in \mathbb{O}$ and A does not depend on x and D_x (the concept of Gröbner bases¹⁶ plays a crucial rôle in this step). Then after integration we get,

$$P \int_a^b f dx + \left[Qf \right]_a^b = 0,$$

in other words, we found a (possibly inhomogeneous) equation for the integral in question. The examples below will demonstrate this methodology clearly; we start with the simplest one, the sequence $f_3(n)$.

Lemma 5. *The sequence $f_3(n)$ given in (16), satisfies the P -finite recurrence*

$$\begin{aligned} & (4n+11)(4n+9)(n+1)^3(n+3)^2 f_3(n+2) - \\ & (2n+3)(16n^6 + 144n^5 + 515n^4 + 930n^3 + 888n^2 + 423n + 81) f_3(n+1) + \\ & (4n+3)(4n+1)(n+2)^3 n^2 f_3(n) = 0. \end{aligned}$$

subject to the initial conditions

$$f_3(1) = -\frac{8}{105}, \quad f_3(2) = -\frac{8}{385}.$$

Proof. It is an easy exercise to compute the first values of $f_3(n)$ explicitly with a computer algebra system. Thus we basically have to derive the recurrence. For this purpose, we compute an annihilating ideal I for $(t^2 - 1)^3(Q_n + \epsilon_n \alpha P_n)^4$ which is the expression in the residue (16). For this purpose we apply holonomic closure properties (note that $Q_n + \epsilon_n \alpha P_n$ satisfies the same equations as Q_n itself). The resulting Gröbner basis is too large to be printed here, namely a full page of equations approximately. It is represented in the Ore algebra $\mathbb{C}(n, t)\langle S_n, D_t \rangle$. In the next step we make use of a special algorithm¹⁷ for computing a creative telescoping operator

$$A(n, S_n) + D_t B(n, t, S_n, D_t) \in I$$

(its existence is guaranteed by the theory of holonomy). Because we are dealing with a residue we can forget about the part B and find that A annihilates the residue. In order to obtain $f_3(n)$ we need to multiply the residue with $2\epsilon_n^{-2}$, which can be done again by closure properties. The resulting operator represents exactly the above recurrence. All these computations were done with the above mentioned package `HolonomicFunctions`^{11,12}. \square

Lemma 6. *The sequence $f_1(n)$ given in (14) satisfies the P-finite recurrence*

$$\begin{aligned} & (4n + 11)(4n + 9)(n + 4)^2(n + 1)^3(4n^2 + 8n - 9)f_1(n + 2) - \\ & (2n + 3)(64n^8 + 768n^7 + 3580n^6 + 8028n^5 + 8113n^4 + \\ & \quad 834n^3 - 4863n^2 - 3276n - 648)f_1(n + 1) + \\ & (4n + 3)(4n + 1)(n + 2)^3(n - 1)^2(4n^2 + 16n + 3)f_1(n) = 0 \end{aligned}$$

subject to the initial conditions

$$f_1(2) = \frac{16}{1155}, \quad f_1(3) = \frac{16}{2145}.$$

Proof. The proof is based on the same ideas as in Lemma 5, except that the expression of which we have to take the residue is more complicated. In particular, an indefinite integral occurs (recall that indefinite integration is not among the holonomic closure properties) and it is not clear a priori how to choose the integration constant such that the result is again holonomic. We start by computing an annihilating ideal I for

$$F(n, t) = (t^2 - 1)^2(Q_n + \epsilon_n \alpha P_n)^2.$$

Thus for all $A \in I$ the operator AD_t annihilates the indefinite integral $\int F(n, t) dt$. Additionally, from a creative telescoping operator $A + D_t B \in I$ we can derive more such annihilating operators. Let J denote the annihilating ideal for $B(F)$ which can be obtained by holonomic closure properties. Then for every $C \in J$, the operator CA annihilates the indefinite integral as well. Altogether we obtain a zero-dimensional annihilating ideal for $\int F(n, t) dt$, and continue as in Lemma 5. \square

These recurrences in Lemmas 5 and 6 are irreducible (in the sense that the corresponding operator cannot be factorized), and so we are not able to find closed forms for f_1 and f_3 . The recurrence for $f_2(2n)$ is given by a third-order recurrence with polynomial coefficients of degree larger than 50, which we do not state here explicitly. The initial conditions are

$$f_2(2) = \frac{16}{1155}, \quad f_2(4) = \frac{184}{183141}, \quad f_2(6) = \frac{38308}{181081875}.$$

This recurrence is reducible and possesses a hypergeometric solution

$$f_2(2) \frac{8\pi^2 \Gamma(n+1) \Gamma(5/6+n)^2 \Gamma(1/6+n)^2 \Gamma(n)^3}{25 \Gamma(n+2/3)^2 \Gamma(3/2+n)^3 \Gamma(1/2+n) \Gamma(4/3+n)^2}$$

but because $f_2(2) \neq 0$, the recurrence for $f_2(2n)$ cannot be reduced.

We are interested in a **practical** way to apply the third-order variational equation. To do this, these recurrences are not enough, since we need closed forms. As these closed forms do not exist, we will instead produce closed form expressions which approach f_1 , f_2 , and f_3 with a controlled relative error. In the following, we will denote the harmonic numbers by

$$H(n) = \sum_{i=1}^{n-1} \frac{1}{i}.$$

Definition 6. *Let us consider an operator $L \in \mathbb{C}\langle n, S_n \rangle$, in other words L represents a linear recurrence with polynomial coefficients. We will say that L is regular at infinity if for all solutions u (i.e., $Lu = 0$) there exist $\alpha \in \mathbb{Z}$, $\beta \in \mathbb{N}$, and $\gamma \in \mathbb{C}$ such that*

$$u(n) \sim \gamma n^\alpha H(n)^\beta \quad \text{for } n \rightarrow \infty.$$

Theorem 7. *Consider $L \in \mathbb{C}\langle n, S_n \rangle$ of order k and assume that it is regular at infinity. Then for all $p \in \mathbb{N}$ and for all u solution of $Lu = 0$, there exists a function $F \in \mathbb{C}(n)[H(n)]$ with degree in $H(n)$ less than $k - 1$ such that*

$$u(n) = F(n) + O\left(\frac{H(n)^{k-1}}{n^p}\right).$$

This theorem is directly implied by the theorem of Birkoff given in¹⁸, which gives a general form of an asymptotic expansion which is always possible. In our case, we will only use what we call the regular case, which in a Birkoff expansion corresponds to not having an exponential part.

Definition 7. Consider a function $f : \mathbb{N} \rightarrow \mathbb{R}$ and a function $F \in \mathbb{R}(n)[H(n)]$. We say that F is an approximation of f with relative error ϵ at rank n_0 if

$$\left| \frac{f(n)}{F(n)} - 1 \right| \leq \epsilon \quad \forall n \geq n_0.$$

We consider p functions $f_1, \dots, f_p : \mathbb{N} \rightarrow \mathbb{R}$ and approximations $F_1, \dots, F_p \in \mathbb{R}(n)[H(n)]$ with relative error ϵ at rank n_0 . We define the error amplification factor A by

$$A = \min \left\{ \tilde{A} \in \mathbb{R}_+^* \text{ such that } \left| \frac{\sum_{i=1}^p f_i(n)}{\sum_{i=1}^p F_i(n)} - 1 \right| \leq \tilde{A}\epsilon \quad \forall n \geq n_0 \right\}.$$

Lemma 7. We consider p functions $f_1, \dots, f_p : \mathbb{N} \rightarrow \mathbb{R}$ and approximations $F_1, \dots, F_p \in \mathbb{R}(n)[H(n)]$ with relative error $\epsilon < 1$ at rank n_0 and A their amplification factor. Then

$$A \leq \max_{n \geq n_0} \frac{\sum_{i=1}^p |F_i(n)|}{\left| \sum_{i=1}^p F_i(n) \right|}.$$

Proof. The lemma is equivalent to prove that

$$\left| \frac{\sum_{i=1}^p f_i(n)}{\sum_{i=1}^p F_i(n)} - 1 \right| \leq \epsilon \max_{n \geq n_0} \frac{\sum_{i=1}^p |F_i(n)|}{\left| \sum_{i=1}^p F_i(n) \right|}$$

So one just needs to maximize the left hand side. We already know that $|f_i(n)/F_i(n) - 1| \leq \epsilon$. So depending on the sign of $f_i(n)$ we replace $f_i(n)$ by $(1 - \epsilon)F_i(n)$ or $(1 + \epsilon)F_i(n)$. We then expand

$$\begin{aligned} \left| \frac{\sum_{i=1}^p f_i(n)}{\sum_{i=1}^p F_i(n)} - 1 \right| &\leq \left| \frac{\epsilon \sum_{i=1}^p \text{sign}(f_i(n)) F_i(n)}{\sum_{i=1}^p F_i(n)} \right| \leq \left| \frac{\epsilon \sum_{i=1}^p |F_i(n)|}{\sum_{i=1}^p F_i(n)} \right| \\ &\leq \epsilon \max_{n \geq n_0} \frac{\sum_{i=1}^p |F_i(n)|}{\left| \sum_{i=1}^p F_i(n) \right|} \end{aligned}$$

using the fact that $f_i(n)$ and $F_i(n)$ have always the same sign for $n \geq n_0$ (because $\epsilon < 1$). \square

Integrability conditions for homogeneous potentials

In practice, we first check that the sign of the functions $F_i(n)$ and their sum does not change for $n \geq n_0$ and then we prove a majoration of the resulting expression in $\mathbb{R}(n, H(n))$. So all comes down to prove that some polynomial in $\mathbb{R}[n, H(n)]$ does not vanish for $n \geq n_0$. This can be done by first making an encadrement of the function $H(n)$ and then prove that the corresponding bivariate polynomial does not vanish on a particular algebraic subset. Such a problem can be algorithmically decided.

Theorem 8. *Consider the recurrence equation*

$$u(n+1) = A(n)u(n) \quad \forall n \in \mathbb{N}, \quad A(n) \in M_p(\mathbb{C}) \quad (17)$$

Consider $\|\cdot\|$ a matricial norm and $R(n)$ the resolvent matrix of equation (17). Assume that

$$M(\infty) = \sum_{j=0}^{\infty} \|A(j) - I_p\| < 1.$$

Then

$$\|R(n) - I_p\| \leq \frac{M(\infty)}{1 - M(\infty)} \quad \forall n \in \mathbb{N}.$$

Proof. We write

$$R(n) = \prod_{i=0}^{n-1} A(i) = \prod_{i=0}^{n-1} ((A(i) - I_p) + I_p).$$

Let us pose

$$M(n) = \sum_{j=0}^{n-1} \|A(j) - I_p\|.$$

We want to prove a majoration of the type

$$\|R(n) - I_p\| \leq CM(n) \quad (18)$$

with a suitable constant $C > 0$. For $n = 1$, this is true with $C = 1$. Let us prove equation (18) by recurrence:

$$R(j) = \prod_{i=0}^{j-1} ((A(i) - I_p) + I_p) = (A(j-1) - I_p) \prod_{i=0}^{j-2} A(i) + \prod_{i=0}^{j-2} A(i),$$

$$R(j) - R(j-1) = (A(j-1) - I_p) \prod_{i=0}^{j-2} A(i) = (A(j-1) - I_p)R(j-1).$$

Then we sum these equations for $1 \leq j \leq n$ which produces

$$\begin{aligned} \|R(n) - I_p\| &= \left\| \sum_{j=0}^{n-1} (A(j) - I_p)(R(j) - I_p) + (A(j) - I_p) \right\| \\ &\leq \sum_{j=0}^{n-1} \|A(j) - I_p\| \|R(j) - I_p\| + \|A(j) - I_p\| \\ &\leq M(n) + \sum_{j=0}^{n-1} \|A(j) - I_p\| CM(j) = M(n) + CM(n)^2 \\ &\leq (1 + CM(\infty))M(n) \end{aligned}$$

using the fact that $M(n)$ is a growing sequence. So the recurrence property is proved if $C \leq 1 + CM(\infty)$ which is equivalent to $C \geq (1 - M(\infty))^{-1} \geq 1$. So this proves that

$$\|R(n) - I_p\| \leq \frac{M(n)}{1 - M(\infty)} \leq \frac{M(\infty)}{1 - M(\infty)}$$

which proves the theorem. □

The main application of this theorem is to compute a sequence with controlled error. Let us take an operator $L \in \mathbb{R}\langle n, S_n \rangle$ regular at infinity. We can then compute an asymptotic expansion of the resolvent matrix of L , and an error matrix which will satisfy an equation like (17). Then for an $n_0 \in \mathbb{N}$, we can apply Theorem 8 for the shifted sequence $u(n + n_0)$, and the majoration $M(\infty)$ will become very small for n_0 big enough, giving us that the error is always lower than some explicit bound. This has very important consequences for the application of the higher variational method. In particular, it becomes possible to rigorously prove that a sequence of potentials with the unbounded eigenvalue property does not satisfy integrability criteria for λ large enough, and thus coming back to a bounded eigenvalue problem.

VII. APPLICATION AT ORDER 2

We now apply the second order criterion to our example. We begin with the case E_4 . Before we state the corresponding theorem, we need a preparatory lemma concerning the solutions of a certain Diophantine equation.

Lemma 8. *The set of solutions $(k_1, k_2) \in \mathbb{N}^2$ of the Diophantine equation*

$$R(k_1, k_2) = k_2^2 k_1^2 + k_2 k_1^2 - 75 k_1^2 - 75 k_1 + k_2 k_1 - 27 k_2 + k_2^2 k_1 - 27 k_2^2 = 0$$

is given by $\{(0, 0), (6, 14)\}$.

Proof. We begin by proving that for $k_2 \geq 50$, the condition $R = 0$ implies $4 < k_1 < 5$, and similarly, for $k_1 \geq 50$, we have $8 < k_2 < 9$. These statements can be written as logical expressions involving polynomial inequalities

$$\forall k_1 \forall k_2 : (k_1 \geq 0 \wedge k_2 \geq 50 \wedge R(k_1, k_2) = 0) \implies 4 < k_1 < 5, \quad (19)$$

$$\forall k_1 \forall k_2 : (k_1 \geq 50 \wedge k_2 \geq 0 \wedge R(k_1, k_2) = 0) \implies 8 < k_2 < 9. \quad (20)$$

Such formulas can be proven routinely with quantifier elimination techniques like cylindrical algebraic decomposition²⁰. Indeed, applying the Mathematica command **CylindricalDecomposition** to the above formulae reveals that they are true. Therefore, there are no integer solutions for $k_1 \geq 50$ or $k_2 \geq 50$ and an exhaustive search delivers exactly the solutions claimed above.

However, if we want to prove (19) and (20) “by hand” (let’s consider the first one for the moment), we have to look at the largest real root of the polynomial

$$\text{res}_{k_1} \left(R(k_1, k_2), \frac{\partial R(k_1, k_2)}{\partial k_1} \right) R(4, k_2) R(5, k_2).$$

We find that this root is smaller than 50 (using real root isolation) and that the limit

$$\lim_{k_2 \rightarrow \infty} \kappa(k_2) = -\frac{1}{2} + \frac{1}{2}\sqrt{109}$$

is between 4 and 5, where $\kappa(k_2)$ denotes the positive solution of $R(k_1, k_2) = 0$ regarded as an equation in k_1 . The implication (19) follows, and (20) can be proven analogously. \square

Theorem 9. *We consider the potential E_4 given in Theorem 5. If the variational equation near all Darboux points is integrable at order 2, then the corresponding eigenvalues are integers of the form $\lambda = (2l - 1)(l + 1)$, $l \in \mathbb{N}$.*

Proof. We use the notation $U = rE_4$ from Theorem 5. The condition $U'(\theta) = 0$ yields the two Darboux points

$$\begin{aligned} c_1 : e^{i\theta} &= 1, \\ c_2 : e^{i\theta} &= \frac{s + 6\lambda_1}{s - 6\lambda_2}. \end{aligned} \quad (21)$$

There are singular cases of the second equation, namely for $s + 6\lambda_1 = 0$ or $s - 6\lambda_2 = 0$. After solving and replacing, we find that these cases correspond exactly to $k_1 = 0$ and $k_1 = 3$, which were excluded from E_4 .

We now compute the third derivative of V , evaluated at the two Darboux points c_1 and c_2 given by expression (21):

$$\begin{aligned}\frac{\partial^3 V}{\partial q_2^3}(c_1) &= \frac{i\lambda_1(s + 15\lambda_1 + 9\lambda_2)}{\lambda_1 + \lambda_2}, \\ \frac{\partial^3 V}{\partial q_2^3}(c_2) &= -\frac{i\lambda_2(s - 15\lambda_2 - 9\lambda_1)}{3(\lambda_1 + \lambda_2)}.\end{aligned}$$

In the case (k_1, k_2) both odd, both derivatives should vanish. We solve the system and we find $4i(k_2 + 1)k_2 = 0$. This is impossible for odd values. In the case k_1 odd k_2 even, the first one should vanish, and in the case k_1 even k_2 odd the second one should vanish. We get the equations

$$\begin{aligned}\frac{k_1^2(k_1 + 1)^2(k_2^2k_1^2 + k_2^2k_1 - 27k_2^2 - 27k_2 - 75k_1 + k_2k_1 - 75k_1^2 + k_2k_1^2)}{12(k_2^2 + k_2 + k_1 + k_2^2)} \\ \frac{k_2^2(k_2 + 1)^2(k_1^2k_2^2 + k_1^2k_2 - 27k_1^2 - 27k_1 - 75k_2 + k_1k_2 - 75k_2^2 + k_1k_2^2)}{12(k_1^2 + k_1 + k_2 + k_2^2)}\end{aligned}\tag{22}$$

These two conditions are symmetric. The first terms can never vanish because we have k_1 odd for the first one and k_2 odd for the second one. To conclude, we need to look at the last term, which corresponds to a Diophantine equation, and to prove that this equation does not have a solution with k_1 odd and k_2 even.

With Lemma 8, we have no solutions from the second term where k_1 and k_2 have different parity. We conclude that all the possibilities left are for k_1, k_2 even. \square

It is well known that Diophantine equations in general cannot be solved (Matiyasevich's theorem). This means that Lemma 8 is a lucky case, although not trivial to prove. We therefore should remark that the study of this equation is not absolutely mandatory. We could simply skip it, **assume** that it is satisfied and continue further to the third-order condition. This condition would add two additional equations in k_1 and k_2 and thus would allow to solve the problem in all generality.

Here we are in a special case. A Diophantine equation $R(k_1, k_2) = 0$ can be solved only using real algebraic geometry in one of the following cases:

1. The set $R^{-1}(0) \cap \mathbb{R}^{+2}$ is compact. In this case we only have a finite number of points to test.
2. The set $R^{-1}(0) \cap \mathbb{R}^{+2}$ is not compact but all infinite branches are asymptotes and the corresponding asymptotic straight lines have a rational slope. In this case, either R

is homogeneous and has an infinite number of solutions, or the integer solutions can be bounded: when approaching infinity, the infinite branch of $R^{-1}(0)$ comes closer to the asymptotic line without touching it; for rational slope, there is then a nonzero infimum for the distance between the asymptotic straight line and integer points).

The first case can be considered to be part of the second one with no asymptotes at all. In Lemma 8, we encounter the second case.

Remark 4. *The potential corresponding to $k_1, k_2 = (6, 14)$ is the following (with the good choice of valuation for the square root):*

$$V(r, \theta) = \frac{1}{r} \left(-20 + \frac{105}{2}e^{i\theta} - 42e^{2i\theta} + \frac{21}{2}e^{3i\theta} \right).$$

This potential has two Darboux points, it is integrable at order 2 near these two Darboux points and we have also that the third derivative near one of the Darboux points is zero (which is not needed for integrability at order 2 but gives interesting properties in practice at order 3).

Theorem 10. *Among the potentials in the families E_1, E_2, E_3 , if a potential V is meromorphically integrable, then it is of the form (after multiplying by some constant factor):*

$$\begin{aligned} V &= \frac{1}{r} \left(-\frac{1}{3}k(2k+1)e^{3i\theta} + \frac{1}{2}k(2k+1)e^{2i\theta} - \frac{1}{6}(2k^2+k-6) \right), \\ V &= \frac{1}{r} \left(-\frac{1}{2}k(2k+1)e^{2i\theta} + k(2k+1)e^{i\theta} - \frac{1}{2}(2k^2+k-2) \right), \end{aligned}$$

for $k \in \mathbb{N}$.

Proof. The potentials E_2 and E_3 possess only one Darboux point. The corresponding potentials are

$$\begin{aligned} E_2 : V &= r^{-1} \left(-\frac{1}{6}k(k+1)e^{3i\theta} + \frac{1}{4}k(k+1)e^{2i\theta} - \frac{1}{12}k^2 - \frac{1}{12}k + 1 \right), \\ E_3 : V &= r^{-1} \left(-\frac{1}{4}k(k+1)e^{2i\theta} + \frac{1}{2}k(k+1)e^{i\theta} - \frac{1}{4}k^2 - \frac{1}{4}k + 1 \right). \end{aligned}$$

We know that if k is odd, we have an additional integrability condition at order 2. We find that

$$\frac{\partial^3 V}{\partial q_2^3}(c) = \frac{5}{2}ik(k+1) \text{ for } E_2, \quad \frac{\partial^3 V}{\partial q_2^3}(c) = \frac{3}{2}ik(k+1) \text{ for } E_3.$$

These terms should vanish. This is never fulfilled for odd k . The sequence of potentials given by Theorem 10 corresponds exactly to the cases of even k (for which there is no condition for

Integrability conditions for homogeneous potentials

integrability at order 2). At last, we have the potential E_1 . The corresponding eigenvalue is always -1 , so it is always integrable at order 2. At order 3, we know that the integrability condition is $U^{(3)}(0) = 0$. We get

$$U^{(3)}(0) = -2ib$$

So the only possibility is $b = 0$ and this corresponds to the potential $V = r^{-1}$. This potential is integrable and already belongs to the family described by Theorem 10.

□

VIII. APPLICATION AT ORDER 3

We will now prove Theorem 3, building an algorithm to prove it.

Proof. The scheme of the proof is the following

- First we prove that the recurrences for f_1, f_2, f_3 are regular at infinity.
- We then produce a series expansion $\tilde{R}_i(n)$ at infinity at an order high enough of the resolvent matrix $R_i(n)$ associated to these recurrences.
- We then write $R_i(n) = \tilde{R}_i(n)\tilde{R}_i(n_0)^{-1}R_i(n_0)E_i(n)$ for a large enough $n_0 \in \mathbb{N}$ and build a recurrence of the form (17) whose resolvent matrix is $E_i(n)$ (after change of basis), which will be denoted by $E_i(n+1) = A_i(n)E_i(n)$. We have moreover that $E_i(n_0)$ is the identity matrix.
- As $\tilde{R}_i(n)$ is a good approximation of $R_i(n)$ when $n \rightarrow \infty$, the matrix $A_i(n)$ will tend to the identity matrix when $n \rightarrow \infty$. Using Theorem 8 with a shift in the indices, we will have that

$$\|E_i(n) - I\| \leq \frac{\sum_{j=n_0}^{\infty} \|A_i(j) - I\|}{1 - \sum_{j=n_0}^{\infty} \|A_i(j) - I\|} \quad \forall n \geq n_0$$

- If we have chosen an expansion order and n_0 large enough, this sum will be finite and small, and thus will give us an approximation of $R_i(n)$ by $\tilde{R}_i(n)$ with relative error control. The expressions in Theorem 3 follow.

Integrability conditions for homogeneous potentials

For $f_3(2n)$, we find the following asymptotic expansion (a high order makes up the computation easier for error control)

$$c_1 \left(\frac{1}{n^4} - \frac{1}{n^5} + \frac{25}{32n^6} - \frac{35}{64n^7} + \frac{183}{512n^8} \right) + c_2 \left(\left(\frac{3}{16n^4} - \frac{3}{16n^5} + \frac{75}{512n^6} - \frac{105}{1024n^7} + \frac{549}{8192n^8} \right) H(n) + \frac{1}{n^2} - \frac{1}{2n^3} + \frac{19951}{46848n^4} - \frac{7507}{46848n^5} + \frac{96541}{1499136n^6} - \frac{58151}{2998272n^7} \right)$$

This proves by the way that the recurrence for $f_3(2n)$ is regular. We do the same for $f_1(2n)$ and $f_2(2n)$ and we find that they are regular too. We then find a majoration of the norm of the error matrix $A_3(n)$

$$\|A_3(n)\|_\infty \leq \frac{9975}{256n^6} + \frac{29925}{4096} \frac{H(n)}{n^6} + \frac{9975}{256n^8} + \frac{29925}{4096} \frac{H(n)}{n^8}$$

We choose now $n_0 = 100$. We majorate the sum of this majoration beginning at $n = 100$.

We find a majoration of this sum by

$$\sum_{n=100}^{\infty} \|A_3(n)\|_\infty \leq 4.84522 \times 10^{-9}$$

$$\|E_3(n)\| \leq \frac{4.84522 \times 10^{-9}}{1 - 4.84522 \times 10^{-9}} \quad \forall n \geq n_0$$

(an explicit rational number). We then compute the recurrence up to $n = 100$, and then produce an encadrement (with error less than 10^{-10}) of the result with rational numbers. Although it is not mandatory in theory, in practice recurrences tend to produce very large rational numbers, whose size grows linearly with n , and thus are impractical to manipulate. This gives us the coefficients c_1, c_2 with a good error control:

$$c_1 = -\frac{883919839}{274877906944}, \quad c_2 = -\frac{1740684681}{8589934592}.$$

We then compute the error amplification of the sum, and find that it is less than $33/32$. As the resulting expression is too complicated to manipulate for applications, we only keep the terms up to order 3 and prove that this new approximation has a relative error less than 10^{-5} . The expressions for f_1 and f_2 are found with a similar way, with the exception that at the end, to produce a sufficiently simple and accurate formula, it is not sufficient to keep the terms up to order 2 (after there is a $H(n)$ that we want to avoid), so we need to add a term of order 3 (without $H(n)$) with a well chosen coefficient such that the error stays below 10^{-5} (else the result is only accurate to 10^{-3}). \square

Theorem 11. *The third order integrability conditions for the families*

$$V = \frac{1}{r} \left(-\frac{1}{3}k(2k+1)e^{3i\theta} + \frac{1}{2}k(2k+1)e^{2i\theta} - \frac{1}{6}(2k^2+k-6) \right)$$

$$V = \frac{1}{r} \left(-\frac{1}{2}k(2k+1)e^{2i\theta} + k(2k+1)e^{i\theta} - \frac{1}{2}(2k^2+k-2) \right)$$

where $k \in \mathbb{N}^*$, are

$$9(k+1)^2(2k-1)^2f_1(2k) = 25k^2(2k+1)^2f_2(2k) + (66k^2+33k-9)f_3(2k),$$

$$9(k+1)^2(2k-1)^2f_1(2k) = 9k^2(2k+1)^2f_2(2k) + (42k^2+21k-9)f_3(2k),$$

respectively. They are never satisfied.

Proof. We replace $f_1(2k), f_2(2k), f_3(2k)$ by their approximations, and then compute the error amplification. It is less than $33/32$, and the resulting expression does not vanish for $k \geq 100$. For $k < 100$, we make exhaustive testing and we do not find any solutions. For the second equation, we do not find any solution either. \square

Theorem 12. *We consider the family of potentials E_4*

$$E_4 : \quad V = r^{-1} \left(\frac{(s-6\lambda_2)\lambda_2}{18(\lambda_1+\lambda_2)} e^{3i\theta} - \frac{(3\lambda_1+s-3\lambda_2)\lambda_2}{6(\lambda_1+\lambda_2)} e^{2i\theta} + \frac{(6\lambda_1+s)\lambda_2}{6(\lambda_1+\lambda_2)} e^{i\theta} + \frac{-9\lambda_1\lambda_2 - \lambda_2s + 18\lambda_1 + 18\lambda_2 - 3\lambda_2^2}{18(\lambda_1+\lambda_2)} \right)$$

with

$$s^2 = 6\lambda_1^2\lambda_2 + 6\lambda_1\lambda_2^2 - 36\lambda_1\lambda_2 \quad \lambda_1 = \frac{1}{2}(k_1-1)(k_1+2) + 1$$

$$\lambda_2 = \frac{1}{2}(k_2-1)(k_2+2) + 1 \quad k_1, k_2 \in \mathbb{N}^* \quad k_1 \neq 3$$

The third order integrability condition for E_4 is of the form

$$Q_{k_1, k_2}(f_1(k_1), f_2(k_1), f_3(k_1)) = 0$$

$$Q_{k_2, k_1}(f_1(k_2), f_2(k_2), f_3(k_2)) = 0$$

where Q is a quadratic form depending polynomially on k_1 and k_2 .

Proof. We use Theorem 2 and compute the derivatives of the potentials in the family E_4 . These derivatives depend rationally on k_1, k_2 , and s . As there are two Darboux points, we get two conditions $(C_1), (C_2)$ linearly dependent on $f_1(k_1), f_2(k_1), f_3(k_1)$ or $f_1(k_2), f_2(k_2), f_3(k_2)$

Integrability conditions for homogeneous potentials

respectively for each Darboux point. To remove the quadratic extension s , we make the product $(C_1) \times \text{subs}(s = -s, (C_1))$ and $(C_2) \times \text{subs}(s = -s, (C_2))$. The fact that in the potentials of E_4 , the two parameters λ_1 and λ_2 play a symmetric rôle produces the two conditions $Q_{k_1, k_2} = 0$ and $Q_{k_2, k_1} = 0$. \square

Remark 5. *Indeed, the conditions $Q_{k_1, k_2}, Q_{k_2, k_1}$ are not equivalent to the conditions $(C_1), (C_2)$. We can solve (C_1) in the quadratic extension and get for example that s should be rational because f_1, f_2, f_3 are always rational (this can be proven even without the P -finite recurrences since they correspond to a particular term in the series expansion of rational expressions in $t, P_n(t), Q_n(t)$). We get that*

$$\sqrt{3k_1k_2(k_2 + 1)(k_1 + 1)(k_1 + k_1^2 + k_2 + k_2^2 - 12)} \in \mathbb{N} \quad (23)$$

if some generic condition depending on the $f_i(k_2), f_i(k_1)$ is satisfied. It corresponds to a Diophantine equation but it does not possess the nice properties we used to solve Lemma 8. We know moreover that (k_1, k_2) should be even. A direct search produces the picture given in Figure ??.

Theorem 13. *The third order integrability condition for E_4 is never satisfied except for $(k_1, k_2) = (2, 2)$.*

Proof. Recall that the parameters (k_1, k_2) need to be both even for a potential E_4 to be integrable at order 2 near all Darboux points. We begin by solving $Q_{k_2, k_1}(f_1(k_2), f_2(k_2), f_3(k_2)) = 0$ in k_1 . This is a polynomial of degree 4 in k_1 and as a polynomial, its Galois group is D_4 . This allows us to write the solution in a relatively simple form

$$k_1 = -\frac{1}{2} + \sqrt{F_1(k_2) + wF_2(k_2)} \quad \text{with} \quad w^2 = 9(k_2 + 2)^2(k_2 - 1)^2 f_1(k_2)f_2(k_2) - 6(k_2 + 3)(k_2 - 2)f_2(k_2)f_3(k_2) + 36f_3(k_2)^2 \quad (24)$$

where $F_1, F_2 \in \mathbb{Q}(f_1, f_2, f_3, k_2)$. Moreover, k_1, k_2 are even integers. Let us prove that in fact, for even $k_2 \geq 200$, the expression

$$-\frac{1}{2} + \sqrt{F_1(k_2) + wF_2(k_2)}$$

is always complex for all possible valuations of the square roots. To have real values, we need that $F_1(k_2) + wF_2(k_2)$ be positive for at least one valuation of the square root. Let us

begin by proving that w never vanishes. The function w^2 is a polynomial in $\mathbb{Q}[f_1, f_2, f_3, k_2]$. Thanks to Theorem 3, we can express f_1, f_2, f_3 in k_2 with controlled relative error. We check that the amplification of the error is small after summation of all terms (here it is less than $1 + 10^{-3}$) and that the approximated expressions never vanish. Now we need to prove that

$$F_1(k_2) + wF_2(k_2) < 0 \text{ and } F_1(k_2) - wF_2(k_2) < 0.$$

We first prove that $F_2(k_2)$ and $F_1(k_2)$ (which are in $\mathbb{Q}[f_1, f_2, f_3, k_2]$ of degree 3, 4 in f_i respectively) are always negative. Then we just have to prove that

$$\frac{F_1(k_2)}{wF_2(k_2)} > 1 \iff \frac{F_1(k_2)^2}{w^2F_2(k_2)^2} > 1 \iff F_1(k_2)^2 - w^2F_2(k_2)^2 > 0.$$

The last expression is in $\mathbb{Q}[f_1, f_2, f_3, k_2]$ (of degree 8 in f_i), so we can prove this statement. Again we compute the error amplification of the sum and it stays below $1 + 10^{-3}$, and the error is then still less than 10^{-4} . Eventually, we prove that this approximated expression never vanishes and is always positive. For the remaining cases, we use exhaustive testing and we find only one solution $(k_1, k_2) = (2, 2)$. \square

The case $(k_1, k_2) = (2, 2)$ corresponds to the second case of Theorem 4. It is really integrable with a quadratic in momenta additional first integral which is given in⁷ page 107 case (8).

IX. REMAINING CASES AND CONCLUSION

The remaining cases are the ones which do not possess a non-degenerate Darboux point.

Theorem 14. *Consider the set of potentials V given by (6) and suppose that V does not possess a non-degenerate Darboux point c . If V is meromorphically integrable, then V belongs to one of the families*

$$\begin{aligned} V &= \frac{1}{r} (a + be^{i\theta}), & V &= \frac{1}{r} (a + be^{2i\theta}), \\ V &= \frac{1}{r} (a + be^{3i\theta}), & V &= \frac{1}{r} (a + be^{i\theta})^3, \end{aligned}$$

with $a \in \mathbb{C}$, $b \in \mathbb{C}^*$.

Proof. First let us suppose that V does not possess any Darboux point c . This means that the function

$$U(\theta) = a + be^{i\theta} + ce^{2i\theta} + de^{3i\theta}$$

does not possess any critical point. The only possibility is that $U(\theta) = F(e^{i\theta})$ with $F(z) = a + bz^n$, $b \neq 0$. This corresponds to the three first cases of Theorem 14. Now suppose there exists one Darboux point c but degenerate. After rotation, we can suppose that the Darboux point corresponds to $\theta = 0$. We have moreover the integrability constraint that $U''(0) = 0$. This gives the potential

$$V = \frac{a}{r} (e^{i\theta} - 1)^3.$$

After rotation, this corresponds to the fourth case of Theorem 14. □

The family $V = \frac{1}{r} (a + be^{i\theta})$ is integrable as given in⁷. For the other ones, the integrability status is still unknown. Let us remark now on the open cases. After rotation and dilatation, these cases correspond in fact to a finite number of potentials which are the following:

$$\begin{aligned} V &= r^{-1} e^{2i\theta}, & V &= r^{-1} (e^{2i\theta} - 1), \\ V &= r^{-1} (e^{3i\theta} - 1), & V &= r^{-1} (e^{i\theta} - 1)^3. \end{aligned}$$

We cannot study these cases because we do not have a particular solution to study, or for the last case a sufficiently non-degenerate one (studying degenerate Darboux points with higher variational method is in fact useless and does not give any additional integrability condition). This is of course the main weakness of the Morales-Ramis theory. This is not due to the difficulty of applying the Morales-Ramis theory as we treat it in this article, but much more a fundamental limitation that seems hard to overcome. One approach could consist in looking for special algebraic orbits of these systems using a direct search (following Hietarinta⁷). This is not successful for all these potentials.

To conclude, let us remark that our holonomic approach to higher variational methods is very general, and in no way limited to this example. This could work at least for all problems about integrability of homogeneous potentials, as it allows to compute various higher integrability conditions of any fixed order. This is linked to the fact that the first order variational equation of a natural Hamiltonian system often corresponds to a spectral problem of a second order differential operator, which generates P-finite sequences of functions, which

in turn appear in the study of higher variational equations. We could also wonder if these arbitrary high eigenvalues are really possible, and if this work is only conceptual and in practice useless. Indeed, very high eigenvalues should correspond to very high degree first integrals, and counting the number of conditions and number of free parameters for the existence of such high degree first integrals strongly suggests they do not exist. But this intuition is wrong, as Andrzej J. Maciejewski, Maria Przybylska found quite recently such an example in dimension 3. This is probably linked to the fact that most of integrable cases come from ultra-degenerate cases, as in our analysis: the generic case E_4 contains only one possibility, and when we look at the third order integrability condition, it seems really to be a miracle that this condition could ever be satisfied. On the contrary, the cases without Darboux points contain lots of integrable potentials.

REFERENCES

- ¹J. Morales-Ruiz and J. Ramis, “Galoisian obstructions to integrability of hamiltonian systems,” *Methods and Applications of Analysis*, **8**, 33–96 (2001).
- ²H. Yoshida, “A criterion for the non-existence of an additional integral in Hamiltonian systems with a homogeneous potential,” *Physica D: Nonlinear Phenomena*, **29**, 128–142 (1987).
- ³J. Morales-Ruiz, J. Ramis, and C. Simo, “Integrability of Hamiltonian systems and differential Galois groups of higher variational equations,” *Annales scientifiques de l’Ecole normale supérieure*, **40**, 845–884 (2007).
- ⁴J. Morales-Ruiz and J. Ramis, “Galoisian obstructions to integrability of hamiltonian systems ii,” *Methods and Applications of Analysis*, **8**, 97–112 (2001).
- ⁵T. Combot, “Generic classification of homogeneous potentials of degree -1 in the plane,” arXiv:1110.6130 (2011).
- ⁶T. Kimura, “On Riemann’s equations which are solvable by quadratures,” *Funkcialaj Ekvacioj*, **12**, 269–281 (1970).
- ⁷J. Hietarinta, “A search for integrable two-dimensional Hamiltonian systems with polynomial potential,” *Physics Letters A*, **96**, 273–278 (1983).
- ⁸A. Maciejewski and M. Przybylska, “Darboux points and integrability of Hamiltonian systems with homogeneous polynomial potential,” *Journal of Mathematical Physics*, **46**,

- 062901 (2005).
- ⁹A. Maciejewski and M. Przybylska, “All meromorphically integrable 2D Hamiltonian systems with homogeneous potential of degree 3,” *Physics Letters A*, **327**, 461–473 (2004).
- ¹⁰J. Morales-Ruiz and S. Simon, “On the meromorphic non-integrability of some N -body problems,” *Discrete and Continuous Dynamical Systems (DCDS-A)*, **24**, 1225–1273 (2009).
- ¹¹C. Koutschan, *Advanced Applications of the Holonomic Systems Approach*, Ph.D. thesis, RISC, Johannes Kepler University, Linz, Austria (2009).
- ¹²C. Koutschan, “HolonomicFunctions (User’s Guide),” Tech. Rep. 10-01 (RISC Report Series, Johannes Kepler University Linz, 2010) <http://www.risc.jku.at/research/combinat/HolonomicFunctions/>.
- ¹³T. Combot, “Integrability conditions at order 2 for homogeneous potentials of degree -1 ,” arXiv:1109.5686 (2011).
- ¹⁴A. Aparicio-Monforte and J. Weil, “A reduction method for higher order variational equations of Hamiltonian systems,” in *Symmetries and Related Topics in Differential and Difference Equations: Jairo Charris Seminar 2009, Escuela de Matematicas, Universidad Sergio Arboleda, Bogotá, Colombia*, Contemporary Mathematics, Vol. 549 (American Mathematical Society, 2011).
- ¹⁵D. Zeilberger, “A holonomic systems approach to special functions identities,” *Journal of Computational and Applied Mathematics*, **32**, 321–368 (1990).
- ¹⁶B. Buchberger, *Ein Algorithmus zum Auffinden der Basiselemente des Restklassenrings nach einem nulldimensionalen Polynomideal*, Ph.D. thesis, University of Innsbruck, Innsbruck, Austria (1965).
- ¹⁷C. Koutschan, “A fast approach to creative telescoping,” *Mathematics in Computer Science*, **4**, 259–266 (2010).
- ¹⁸J. Wimp and D. Zeilberger, “Resurrecting the asymptotics of linear recurrences,” *Journal of mathematical analysis and applications*, **111**, 162–176 (1985).
- ¹⁹R. Martínez and C. Simó, “Non-integrability of hamiltonian systems through high order variational equations: Summary of results and examples,” *Regular and Chaotic Dynamics*, **14(3)**, 323–348 (2009).
- ²⁰G. E. Collins, “Quantifier elimination for the elementary theory of real closed fields by cylindrical algebraic decomposition,” *Lecture Notes in Computer Science*, **33**, 134–183 (1975).